arXiv:1508.04450v1 [math.CA] 18 Aug 2015

ON PROBABILISTIC ASPECTS OF CHEBYSHEV POLYNOMIALS

PAWEŁ J. SZABŁOWSKI

Abstract. We expand some, mostly nonnegative rational functions of one or two variables in the series of Chebyshev polynomials. If such a series is multiplied by the density that makes Chebyshev polynomials orthogonal then we obtain Fourier series expansion of certain probabilistic density in the case of rational function of one variable or Lancaster expansion in the case of function of two variables. We study also the general case of rational function of order n depending (symmetrically) on n parameters and find (at least theoretical) expansion.

1. Introduction

The purpose of this note is to recall known and establish new, simple, expansions of some rational functions of one and two variables in the series of Chebyshev polynomials. If such series is multiplied by the density that makes Chebyshev polynomials orthogonal then we would get orthogonal series converging in mean square as well as in almost everywhere sense. Specifically preferred would be those rational functions that would be nonnegative on the segment $[-1,1]$ in the case of functions of one variable or on the square $[-1,1]\times[-1,1]$ in the case of bivariate functions. This would lead to new probability densities on compact sets.

To fix notation we will denote by $\left\{T_n(x)\right\}_{n\geq 0}$ Chebyshev polynomials of the first kind while $\left\{U_n(x)\right\}_{n\geq 0}$ will denote Chebyshev polynomials of the second kind. Recall that these two families satisfy the same 3-term recurrence

$$T_{n+1}(x)=2xT_n(x)-T_{n-1}(x), \tag{1.1}$$

with different initial conditions. Namely $U_0(x)=T_0(x)=1$, $T_1(x)=x$ and $U_1(x)=2x$. Recall also that polynomials T_n are orthogonal with respect to the density $f_C(x)=1/(\pi\sqrt{1-x^2})I_{(-1,1)}$ while polynomials U_n are orthogonal with respect to the density $f_W(x)=\frac{2}{\pi}\sqrt{1-x^2}$ called semicircle or Wigner distribution.

In the sequel we will use exchangeably $\mathbf{a}_n$ and $(\mathbf{a}_1,\ldots,a_n)^T$.

In particular we will analyze the properties of distributions with the following densities:

$$f_{nT}(x|\mathbf{a}_n)=B_n(\mathbf{a}_n)f_C(x)\prod_{j=1}^{n}\varphi_T(x|a_i). \tag{1.2}$$

Here φ_T denotes the generating function of polynomials T_n, i.e. $\varphi_T(x|a)=\sum_{j=0}^{\infty}a^jT_j(x)$ defined for $|a|<1$ and B_n is suitable constant. Obviously $B_1=1$. We will assume

---

Date: August 2015.

2010 *Mathematics Subject Classification.* Primary 41A10,41A50; Secondary 42A16, 62H86.

Key words and phrases. Chebyshev polynomials, bivariate expansions, families of conditional densities, symmetric rational functions, symmetric polynomials.

that $|a_i| < 1$, $i = 1, \ldots, n$. It is well known that

$$\varphi_T(x|a) = (1 - ax)/(1 + a^2 - 2ax). \tag{1.3}$$

It is easy to notice that φ_T is positive for all $|a|, |x| < 1$. Specially interesting will be cases with even n and parameters a_i forming conjugate pairs.

Let us recall that the similar to the density (1.2) case involving Chebyshev polynomials of the second kind was analyzed in [5].

The paper is organized as follows. In the next section we recall some known and obtain some new expansions of rational functions of one and two variables in the series of Chebyshev polynomials. This is done for the sake of completeness of the paper. The Section 3 contains our main results concerning expansion of the density (1.2) in the Fourier series of Chebyshev polynomials of the first kind as well as particular cases for $n = 2, 3, 4$ including cases with parameters a_i forming conjugate pairs. Finally longer proofs are shifted to the last Section 4.

2. AUXILIARY RESULTS

Let us denote for simplicity:

$$w(x, y|\rho) = (1 - \rho^2)^2 - 4xy\rho(1 + \rho^2) + 4\rho^2(x^2 + y^2)$$

and let us recall the so called Poisson–Mehler expansion formula presented and proved in e.g. [1], [2], [6] considered for $q = 0$ gives:

$$g_U(x, y|\rho) \overset{df}{=} \frac{1 - \rho^2}{w(x, y|\rho)} = \sum_{j\geq 0} \rho^j U_j(x) U_j(y), \tag{2.1}$$

convergent everywhere for $|x|, |y| \leq 1$ $|\rho| < 1$ since $\max_{|x|\leq 1} |U_n(x)| = (n + 1)$ and the series $\sum_{j\geq 0} \rho^j (j + 1)^2$ is convergent.

Proposition 1. $\forall |x|, |y| \leq 1, |\rho| < 1:$

$$g_T(x, y|\rho) \overset{df}{=} \frac{1 - \rho^2 - xy\rho(3 + \rho^2) + 2\rho^2(x^2 + y^2)}{w(x, y|\rho)} = \sum_{j=0}^{\infty} \rho^j T_j(x) T_j(y) \geq 0, \tag{2.2}$$

$$g_{UT}(x, y|\rho) \overset{df}{=} \frac{(1 - \rho^2) - 2\rho xy + 2\rho^2 y^2}{w(x, y|\rho)} = \sum_{j=0}^{\infty} \rho^j U_j(x) T_j(y), \tag{2.3}$$

$$\frac{2\rho^2(x - \rho y)}{w(x, y|\rho)} = \sum_{j=0}^{\infty} \rho^j U_{j+1}(x) U_j(y), \tag{2.4}$$

$$\frac{(y - \rho x)(1 + \rho^2 - 2\rho xy)}{w(x, y, \rho)} = \sum_{j=0}^{\infty} \rho^j T_j(x) T_{j+1}(y) \tag{2.5}$$

$$\frac{4x^2 - 1 - \rho}{(1 - \rho)((1 + \rho)^2 - 4\rho x^2)} = \sum_{j=0}^{\infty} \rho^j U_j(x) U_{j+2}(x) \tag{2.6}$$

$$\frac{2(x^2 + y^2 - 2\rho xy)}{w(x, y|\rho)} = x \sum_{n\geq 0}^{\infty} \rho^n U_{n+1}(x) U_n(y) + y \sum_{n\geq 0}^{\infty} \rho^n U_{n+1}(y) U_n(x) \geq 0. \tag{2.7}$$

convergent almost everywhere for $|x|, |y| \leq 1$ $|\rho| < 1$.

Proof. Is shifted to Section 4 □

As a corollary we get the following fact:

Corollary 1. $\forall\, |\rho| < 1,\ |x| \leq 1:$

$$\sum_{j=0}^{\infty} \rho^{2j} T_{2j}(x) = \frac{1+\rho^2-2x^2\rho^2}{(1+\rho^2-2x\rho)(1+\rho^2+2x\rho)} \geq 0, \tag{2.8}$$

$$\sum_{j=0}^{\infty} \rho^{2j+1} T_{2j+1}(x) = \frac{x\rho(1-\rho^2)}{(1+\rho^2-2x\rho)(1+\rho^2+2x\rho)}, \tag{2.9}$$

$$\sum_{j\geq 0} \rho^{j} U_j(x) T_j(x) = \frac{(1+\rho^2-2\rho x^2)}{(1-\rho)((1+\rho)^2-4\rho x^2)} \geq 0, \tag{2.10}$$

$$\sum_{j\geq 0} \rho^{4j} T_{4j}(x) = \frac{1-\rho^4+2\rho^4 x^2(1-x^2)}{((1+\rho^2)^2-2\rho^2x^2)((1-\rho^2)^2+4\rho^2x^2)} \geq 0 \tag{2.11}$$

Proof. (2.8) and (2.9) First we take $x = y$ in (2.2), secondly we use the formula $T_{2n}(x) = 2T_n^2(x) - 1$ and change ρ to ρ^2. (2.10) We put $y = x$ in (2.3). ☐

3. Densities

Theorem 1. *For* $\forall n \geq 1$ *and parameters* $\mathbf{a}_n$ *mutually different we get*

$$B_n(\mathbf{a}_n) = 1/(\frac{1}{2^n} + \frac{1}{2^n}\sum_{k=1}^{n} b_{n,k}),$$

where we denoted $b_{n,k}(\mathbf{a}_n) = \prod_{j=1,j\neq k}^{n} \frac{\gamma(a_k,a_j)}{(a_k-a_j)(1-a_ka_j)}$, $\gamma(a,b) = 2a - b - a^2b$. *Besides we have an expansion:*

$$f_{nT}(\mathbf{a}_n) = f_C(x) + B_n(\mathbf{a}_n) f_C(x) \frac{1}{2^{n-1}} \sum_{j=1}^{\infty} T_j(x) t_j(\mathbf{a}_n),$$

with $t_j(\mathbf{a}_n) = \sum_{k=1}^{n} b_{n,k} a_k^j$.

Proof. Is shifted to Section 4. ☐

To describe briefly constants $B_n(\mathbf{a}_n)$ let us define the following symmetric functions:

$$S_k^{(n)}(\mathbf{a}_n) = \sum_{0\leq j_1<j_2<\ldots,<j_k\leq n} \prod_{m=1}^{m} a_{j_m} \tag{3.1}$$

$$\Delta_m^{(n)}(\mathbf{a}_n) = \sum_{\substack{0\leq j_1,\ldots j_{n-1}\leq m \\ j_1+\ldots+j_{n-1}\leq m}} a_1^{j_1} a_2^{j_2} \ldots a_{n-1}^{j_{n-1}} a_n^{m-(j_1+\ldots+j_{n-1})}, \tag{3.2}$$

$$P_n(\mathbf{a}_n) = \prod_{1\leq i<j\leq n} (1 - a_i a_j). \tag{3.3}$$

As a corollary we have:

Corollary 2. *Using Mathematica we obtain:* $B_2(a_1,a_2) = \frac{2(1-a_1a_2)}{(2-a_1a_2)} = \frac{2P_2(a_1,a_2)}{P_2(a_1,a_2)+1}$,
$B_3(a_1,a_2,a_3) = \frac{4P_3(a_1,a_2,a_3)}{P_2(\mathbf{a}_3)+3-S_2(\mathbf{a}_3)}$,
$B_4(\mathbf{a}_4) = \frac{8P_4(\mathbf{a}_4)}{P_4(\mathbf{a}_4)+7-3S_2(\mathbf{a}_4)+S_3(\mathbf{a}_4)(S_1(\mathbf{a}_4)-S_3(\mathbf{a}_4))-S_4(\mathbf{a}_4)(4-2S_2(\mathbf{a}_4)+3S_4(\mathbf{a}_4))}$.

Let us start with $n = 2$. We have the following observation:

Corollary 3. *Density $f_{2T}(x|a_1,a_2)$ has the expansion:*

$$f_{2T}(x|a_1,a_2)=f_C(x)+B_2f_C(x)\frac{1}{2}\sum_{k=1}^{\infty}T_k(x)(\sum_{j=0}^{k}a_1^ja_2^{k-j}+\frac{a_1^k+a_2^k}{1-a_1a_2}), \tag{3.4}$$

where $B_2=\frac{2-2a_1a_2}{2-a_1a_2}$.

Proof. Following assertions of Theorem 1 all one has to calculate are coefficients $t_j(a_1,a_2)$. Recall that:

$$t_j(a_1,a_2)=\frac{(2a_1-a_2-a_1^2a_2)a_1^j}{(a_1-a_2)(1-a_1a_2)}+\frac{(2a_2-a_1-a_2^2a_1)a_2^j}{(a_2-a_1)(1-a_1a_2)}.$$

Following this we have:

$$\begin{aligned}t_j(a_1,a_2)&=\frac{2(a_1^{j+1}-a_2^{j+1})-a_1a_2(a_1^{j-1}-a_2^{j-1})-a_1a_2(a_1^{j+1}-a_2^{j+1})}{(a_1-a_2)(1-a_1a_2)}\\&=\frac{(2-a_1a_2)}{1-a_1a_2}\sum_{k=0}^{j}a_1^{j-k}a_2^k-\frac{a_1a_2}{1-a_1a_2}\sum_{k=0}^{j-2}a_1^{j-2-k}a_2^k\\&=\sum_{k=0}^{j}a_1^{j-k}a_2^k+\frac{1}{1-a_1a_2}(\sum_{k=0}^{j}a_1^{j-k}a_2^k-\sum_{k=0}^{j-2}a_1^{j-1-k}a_2^{k+1})\\&\sum_{k=0}^{j}a_1^{j-k}a_2^k+\frac{a_1^j+a_2^j}{1-a_1a_2}.\end{aligned}$$

□

3.1. **Complex parameters.** In this subsection we will consider complex parameters a_i, $i=1,\ldots,n$ but assume that n is even and parameters a_i form conjugate pairs. To fix notation let us assume that first two parameters form first conjugate pair, the third and forth the second conjugate pair and so on. Let us denote

$$f_n(x|\rho_1,y_1,\rho_2,y_2,\ldots,\rho_n,y_n)=f_{2nT}(x|\mathbf{a}_{2n}),$$

where we denoted $a_1=\rho_1\exp(i\theta_1),a_2=\rho_1\exp(-i\theta_1),\ldots,a_{2n-1}=\rho_n\exp(i\theta_n),a_{2n}=\rho_n\exp(-i\theta_n))$ and $\theta_i=y_i$, $i=1,\ldots,n$.

Notice that for the pair $a=\rho\exp(i\theta)$, $b=\rho\exp(-i\theta)$ we have $(1+a^2-2ax)(1+b^2-2bx)=w(x,y|\rho)$ and $(1-ax)(1-bx)=1-2\rho xy+\rho^2x^2$, where as before $y=\cos\theta$.

Corollary 4. *$f_2(x|\rho,y)=f_C(x)\frac{2(1-\rho^2)}{2-\rho^2}\frac{1-2xy\rho+\rho^2x^2}{w(x,y|\rho)}$ allows the following expansion:*

$$f_2(x|\rho,y)=f_C(x)+\frac{1-\rho^2}{2-\rho^2}f_C(x)\sum_{j\geq1}\rho^jT_j(x)U_j(y)+\frac{2}{2-\rho^2}f_C(x)\sum_{j\geq1}\rho^jT_j(x)T_j(y). \tag{3.5}$$

Proof. We use (3.4). Next we observe: $\sum_{j=0}^{k}a_1^ja_2^{k-j}=\rho^k\sum_{j=0}^{k}\exp(ij\theta)\exp(-i(k-j)\theta)=\rho^k\exp(-ik\theta)\sum_{j=0}^{k}\exp(2ij\theta)=\rho^k\frac{\exp(-ik\theta)(1-\exp(i(k+1)\theta)}{1-\exp(2i\theta)}=\rho^k\frac{2i\sin(k+1)\theta}{2i\sin\theta}=\rho^kU_k(y)$. Secondly we have: $a_1^k+a_2^k=2\rho^k\cos k\theta=2\rho^kT_k(y)$ and finally; $a_1a_2=\rho^2$. □

Corollary 5. *i)* $\int_{-1}^{1} f_2(x|\rho,y) f_C(y)dy = f_C(x) + \frac{1-\rho^2}{2-\rho^2} f_C(x)\sum_{j\geq 1}\rho^{2j}T_{2j}(x) = f_C(x)\frac{2(1+\rho^2-x^2\rho^2(3-\rho^2))}{(2-\rho^2)((1+\rho^2)^2-4\rho^2x^2)}$.

ii) $\frac{2}{\pi}\int_{-1}^{1} f_2(x|\rho,y)\sqrt{1-y^2}dy = f_C(x)(1-\frac{\rho x}{2-\rho^2})$

Proof. i) The first formula comes straightforwardly from (3.5). The second one comes from application of (2.8).

ii) We utilize the fact that $\forall n\geq 2: \frac{2}{\pi}\int_{-1}^{1} U_n(y)\sqrt{1-y^2}dy = 0$ and $\frac{2}{\pi}\int_{-1}^{1} T_n(y)\sqrt{1-y^2}dy$ $= 0$ while $\frac{2}{\pi}\int_{-1}^{1} U_1(y)\sqrt{1-y^2}dy = 0$ and $\frac{2}{\pi}\int_{-1}^{1} T_1(y)\sqrt{1-y^2}dy = -\frac{1}{2}$. ☐

Remark 1. *Notice that* $f_2(x|y,\rho)$ *as well as the densities exposed in Corollary 5 are conditional densities. Hence in particular for every* $|y|\leq 1$ *and* $|\rho|<1$ *we get* $\int_{-1}^{1} f_2(x|y,\rho)dx = 1$ *or for every* $|\rho|<1$ *we get* $\int_{-1}^{1} f_C(x)\frac{2(1+\rho^2-x^2\rho^2(3-\rho^2))}{(2-\rho^2)((1+\rho^2)^2-4\rho^2x^2)}dx = 1$ *or* $\int_{-1}^{1} f_C(x)(1-\frac{\rho x}{2-\rho^2})dx = 1$. *Besides shapes of* f_2 *for different* y *and* ρ *are very versatile. For example we have plots of* $f_2(x|y,\rho)$ *for* $|x|<1$ *for different values of* y *and* ρ

Here red plot is for $y = .5$, $\rho = .5$, *blue for* $y = -1/5$, $\rho = 4/5$, *green for* $y = 1/4$, $\rho = 3/4$. *One can notice that for some values of* y *and* ρ *the plot has one maximum at some* $x\in(-1,1)$.

Further as $n>2$ is concerned we have:

Proposition 2. *Let* $n = 4$. *For* $a_1 = \rho_1\exp(i\theta_1)$, $a_2 = \rho_1\exp(-i\theta_1)$, $a_3 = \rho_2\exp(i\theta_2)$, $a_4 = \rho_2\exp(-i\theta_2)$ *we get*

$$B_4(\mathbf{a}_4) = B_4(y_1,\rho_1,y_2,\rho_2)$$
$$= \frac{8(1-\rho_1^2)(1-\rho_2^2)w(y_1,y_2,\rho_1\rho_2)}{\alpha(\rho_1,\rho_2)+4(\beta_1(\rho_1,\rho_2)y_1^2+\beta_2(\rho_1,\rho_2)y_2^2)-4\gamma(\rho_1,\rho_2)y_1y_2}$$

where $y_1 = \cos(\theta_1)$, $y_2 = \cos(\theta_2)$, $\alpha(\rho_1,\rho_2) = (1-\rho_1^2\rho_2^2)(4+(4-\rho_1^2\rho_2^2)(1-\rho_1^2)(1-\rho_2^2))$, $\beta_1(\rho_1,\rho_2) = \rho_1^2\rho_2^2(1-\rho_2^2)(2-\rho_1^2)$, $\beta_2(\rho_1,\rho_2) = \rho_1^2\rho_2^2(1-\rho_1^2)(2-\rho_2^2)$, $\gamma(\rho_1,\rho_2) = \rho_1\rho_2((2+\rho_1^2\rho_2^2)(1-\rho_1^2)(1-\rho_2^2)+2(1-\rho_1^2\rho_2^2))$.

Consequently

$$f_4(x|y_1,\rho_1,y_2,\rho_2) = B_4(y_1,\rho_1,y_2,\rho_2)\frac{(1-2\rho_1 xy_1+\rho_1^2x^2)}{w(x,y_1|\rho_1)}\frac{(1-2\rho_2 xy_2+\rho_2^2x^2)}{w(x,y_2|\rho_2)}$$

is a conditional density i.e. $\forall\left|y_1\right|,\left|y_2\right| \leq 1, \left|\rho_1\right|,\left|\rho_2\right| < 1$

$$\int_{\equiv 1}^{1} f_4(x|y_1,\rho_1,y_2,\rho_2)dx = 1.$$

Proof. We have $P_4(\mathbf{a}_4) = (1-\rho_1^2)(1-\rho_2^2)\times(1-\rho_1\rho_2\exp(i(\theta_1-\theta_2))(1-\rho_1\rho_2\exp(-i(\theta_1-\theta_2))\times(1-\rho_1\rho_2\exp(i(\theta_1+\theta_2))(1-\rho_1\rho_2\exp(-i(\theta_1+\theta_2)) = (1-\rho_1^2)(1-\rho_2^2)\times(1+\rho_1^2\rho_2^2-2\rho_1\rho_2\cos(\theta_1-\theta_2))\times(1+\rho_1^2\rho_2^2-2\rho_1\rho_2\cos(\theta_1+\theta_2)) = (1-\rho_1^2)(1-\rho_2^2)((1+\rho_1^2\rho_2^2)^2 - 4y_1y_2\rho_1\rho_2(1+\rho_1^2\rho_2^2)+2\rho_1^2\rho_2^2(2y_1^2-1+2y_2^2-1))$, since $\cos(\theta_1-\theta_2)\cos(\theta_1+\theta_2) = (\cos(2\theta_1)+\cos(2\theta_2))/2$, $\cos(2\theta_1) = T_2(y_1) = 2y_1^2-1$, $\cos(\theta_1-\theta_2)+\cos(\theta_1+\theta_2) = 2y_1y_2$. Consequently $P_4(\mathbf{a}_4) = (1-\rho_1^2)(1-\rho_2^2)w(y_1,y_2,\rho_1\rho_2)$.

$S_4(\mathbf{a}_4) = \rho_1^2\rho_2^2$, $S_3(\mathbf{a}_4) = \rho_1^2\rho_2\exp(i\theta_2)+\rho_1^2\rho_2\exp(-i\theta_2)+\rho_2^2\rho_1\exp(i\theta_1)+\rho_2^2\rho_1\exp(-i\theta_1) = 2\rho_1^2\rho_2y_2+2\rho_2^2\rho_1y_1$,

$S_2(\mathbf{a}_4) = \rho_1^2+\rho_2^2+\rho_1\rho_2(\exp(i(\theta_1+\theta_2)+\exp(-i(\theta_1+\theta_2)+\exp(i(\theta_1-\theta_2)) + \exp(i(\theta_1-\theta_2))) = \rho_1^2+\rho_2^2+\rho_1\rho_2(2\cos(\theta_1+\theta_2)+2\cos(\theta_1-\theta_2)) = \rho_1^2+\rho_2^2 + 4\rho_1\rho_2y_1y_2$, $S_1(\mathbf{a}_4) = 2\rho_1y_1+2\rho_2y_2$. Now we use Corollary 2 and get $P_4(\mathbf{a}_4) + 7-3S_2(\mathbf{a}_4)+S_3(\mathbf{a}_4)(S_1(\mathbf{a}_4)-S_3(\mathbf{a}_4))-S_4(\mathbf{a}_4)(4-2S_2(\mathbf{a}_4)+3S_4(\mathbf{a}_4))) = (1-\rho_1^2)(1-\rho_2)w(y_1,y_2,\rho_1\rho_2)+7-3(\rho_1^2+\rho_2^2+4\rho_1\rho_2y_1y_2)+2\rho_1\rho_2(\rho_1y_2+\rho_2y_1)(2(\rho_1y_1 + \rho_2y_2)-\rho_1^2+\rho_2^2+4\rho_1\rho_2y_1y_2)-\rho_1^2\rho_2^2(4-2(\rho_1^2+\rho_2^2+4\rho_1\rho_2y_1y_2)+3\rho_1^2\rho_2^2)$. □

Remark 2. *As one can check with a help of Mathematica this time for some values of* y_1, y_2, ρ_1, ρ_2 *the plot of* f_4 *has two maxima for some two values of* $x \in (-1,1)$.

4. PROOFS

Proof of the Proposition 1. First let us show (2.3). We will use (2.1) and $T_n(x) = U_n(x) - xU_{n-1}(x)$. We have:

$$\begin{aligned} g_{TU}(x,y|\rho) &= \sum_{j=0}^{\infty}\rho^jU_j(x)T_j(y) = 1+\sum_{j=1}^{\infty}\rho^jU_j(x)(U_j(y)-yU_{j-1}(y)) \\ &= 1+\sum_{j=1}^{\infty}\rho^jU_j(x)U_j(y) - y\sum_{j=1}^{\infty}\rho^jU_j(x)U_{j-1}(y) = \end{aligned}$$

and further

$$\begin{aligned} &= \frac{1-\rho^2}{w(x,y|\rho)} - y\rho\sum_{j=0}^{\infty}\rho^jU_{j+1}(x)U_j(y) = \\ &= \frac{1-\rho^2}{w(x,y|\rho)} - y\rho\sum_{j=0}^{\infty}\rho^j(2xU_j(x)-U_{j-1}(x))U_j(y) = \\ &= \frac{1-\rho^2}{w(x,y|\rho)} - 2xy\rho\frac{1-\rho^2}{w(x,y|\rho)} + y\rho\sum_{j=1}^{\infty}\rho^jU_{j-1}(x)U_j(y) \end{aligned}$$

and finally

$$\begin{aligned}
&= \frac{(1-\rho^2)(1-2xy\rho)}{w(x,y|\rho)} + y\rho^2\sum_{j=0}^{\infty}\rho^j U_j(x)U_{j+1}(y) = \\
&= \frac{(1-\rho^2)(1-2xy\rho)}{w(x,y|\rho)} + y\rho^2\sum_{j=0}^{\infty}\rho^j U_j(x)(2yU_j(y)-U_{j-1}(y)) \\
&= \frac{(1-\rho^2)(1-2xy\rho)}{w(x,y|\rho)} + 2y^2\rho^2\frac{1-\rho^2}{w(x,y|\rho)} - y\rho^2\sum_{j=1}^{\infty}\rho^j U_j(x)U_{j-1}(y)
\end{aligned}$$

Hence we have an equation:

$$\frac{1-\rho^2}{w(x,y|\rho)} - yB = \frac{(1-\rho^2)(1-2xy\rho+2y^2\rho^2)}{w(x,y|\rho)} - y\rho^2 B,$$

where we denoted $B = B(x,y|\rho) = \sum_{j=1}^{\infty}\rho^j U_j(x)U_{j-1}(y)$. So

$$B(x,y|\rho) = \frac{2\rho(x-\rho y)}{w(x,y|\rho)}.$$

and consequently

$$g_{TU}(x,y|\rho) = \frac{(1-\rho^2)}{w(x,y|\rho)} - y\frac{2\rho(x-\rho y)}{w(x,y|\rho)} = \frac{(1-\rho^2)-2\rho xy+2\rho^2y^2}{w(x,y|\rho)}.$$

Hence we have shown (2.3) and (2.4). Using , well known relationship that $T_n(x) = (U_n(x)-U_{n-2}(x))/2$, $n \geq 1$ we have

$$g_T(x,y|\rho) = 1 + \frac{1}{4}\sum_{n\geq 1}\rho^n(U_n(x)-U_{n-2}(x))(U_n(y)-U_{n-2}(y)) \tag{4.1}$$

$$= \frac{3}{4} + \frac{1}{4}g_U(x,y|\rho) + \rho^2 g_U(x,y|\rho)/4 - \frac{1}{4}\sum_{n\geq 1}\rho^n(U_n(x)U_{n-2}(y)+U_{n-2}(x)U_n(y)). \tag{4.2}$$

Now we use formula $U_{n+1}(x) = 2xU_n(x) - U_{n-1}(x)$ getting:

$$g_T(x,y|\rho) = \frac{(1+3\rho^2)}{4}g_U(x,y|\rho) - \frac{1}{4}\rho^2 A(x,y|\rho), \tag{4.3}$$

where $A(x,y|\rho) = x\sum_{n\geq 0}^{\infty}\rho^n U_{n+1}(x)U_n(y) + y\sum_{n\geq 0}^{\infty}\rho^n U_{n+1}(y)U_n(x)$. We calculate $A(x,y|\rho)$ using (2.3) getting

$$A(x,y|\rho) = xB(x,y|\rho)/\rho + yB(y,x|\rho)/\rho = \frac{2(x^2+y^2-2\rho xy)}{w(x,y|\rho)}.$$

Inserting in (4.3) we get (2.2). Thus we have shown (2.2) and (2.7). Now to show (2.5) we have denoting $A(x,y) = \sum_{j=0}^{\infty}\rho^j T_j(x)T_{j+1}(y)$:

$$\begin{aligned}
A(x,y) &= 2y\sum_{j=0}^{\infty}\rho^j T_j(x)T_j(y) - \sum_{j=0}^{\infty}\rho^j T_j(x)T_{j-1}(y) \\
&= 2y\sum_{j=0}^{\infty}\rho^j T_j(x)T_j(y) - y - \rho A(y,x),
\end{aligned}$$

since $T_{-1}(y)=y$. Now iterating we obtain equation:

$$(1-\rho^2)A(x,y)=(y-\rho x)(2\sum_{j=0}^{\infty}\rho^jT_j(x)T_j(y)-1).$$

Now we use (2.2). Finally to get (2.6) we put $y=x$ in equation (4.1, 4.2). ☐

Proof of the Theorem 1. First we will prove expansion

(4.4)

$$\prod_{j=1}^{n}\frac{(1-a_jx)}{(1+a_j^2-2a_jx)}=\frac{1}{2^n}+\frac{1}{2^n}\sum_{k=1}^{n}\prod_{j=1,j\neq k}^{n}\frac{(1-a_k^2)\gamma(a_k,a_j)}{(a_k-a_j)(1-a_ka_j)(1+a_k^2-2a_kx)}.$$

Proof is by induction start with $n=2$ and decomposition of

$$\frac{(1-a_1x)(1-a_2x)}{(1+a_1^2-2a_1x)(1+a_2^2-2a_2x)}=\frac{1}{4}+$$
$$\frac{(1-a_1^2)\gamma(a_1,a_2)}{4(a_1-a_2)(1-a_1a_2)(1+a_1^2-2a_1x)}+\frac{(1-a_2^2)\gamma(a_2,a_1)}{4(a_2-a_1)(1-a_1a_2)(1+a_2^2-2a_2x)},$$

which we get by direct calculation. Now let us assume that (4.4) is true for $n=m$. Hence

$$\prod_{j=1}^{m+1}\frac{(1-a_jx)}{(1+a_j^2-2a_jx)}= \tag{4.5}$$

(4.6)

$$\frac{1-a_{m+1}x}{1+a_{m+1}^2-2a_{m+1}x}(\frac{1}{2^m}+\frac{1}{2^m}\sum_{k=1}^{m}\prod_{j=1,j\neq k}^{m}\frac{(1-a_j^2)\gamma(a_k,a_j)}{(a_k-a_j)(1-a_ka_j)(1+a_k^2-2a_kx)}).$$

Secondly notice that

$$\frac{1-a_{m+1}x}{1+a_{m+1}^2-2a_{m+1}x}=\frac{1}{2}+\frac{(1-a_{m+1}^2)}{2(1+a_{m+1}^2-2a_{m+1}x)}. \tag{4.7}$$

We use this expansion in (4.5) and get

$$\prod_{j=1}^{m+1}\frac{(1-a_jx)}{(1+a_j^2-2a_jx)}=\frac{1}{2^{m+1}}+\frac{1}{2^{m+1}}\sum_{k=1}^{m}\prod_{j=1,j\neq k}^{m}\frac{(1-a_j^2)\gamma(a_k,a_j)}{(a_k-a_j)(1-a_ka_j)(1+a_k^2-2a_kx)}$$
$$+\frac{(1-a_{m+1}^2)}{2^{m+1}(1+a_{m+1}^2-2a_{m+1}x)}$$
$$+\frac{1}{2^{m+1}}\sum_{k=1}^{m}\prod_{j=1,j\neq k}^{m}\frac{(1-a_j^2)\gamma(a_k,a_j)}{(a_k-a_j)(1-a_ka_j)(1+a_k^2-2a_kx)}\frac{(1-a_{m+1}^2)}{2(1+a_{m+1}^2-2a_{m+1}x)}.$$

Now we use expansion:

$$\frac{1}{(1+a_1^2-2a_1x)(1+a_2^2-2a_2x)}=$$
$$\frac{a_1}{(a_1-a_2)(1-a_1a_2)(1+a_1^2-2a_1x)}+\frac{a_2}{(a_2-a_1)(1-a_1a_2)(1+a_2^2-2a_1x)}.$$

applied to (a_1,a_{m+1}), ..., (a_m,a_{m+1}). The next observation is the following:

$$1+\frac{a_k(1-a_{m+1}^2)}{(a_k-a_{m+1})(1-a_ka_{m+1})}=\frac{\gamma(a_k,a_{m+1})}{(a_k-a_{m+1})(1-a_ka_{m+1})}.$$

Hence the sums of coefficients by $1/(1+a_k^2-2a_kx)$ are as the should.

Having proved expansion (4.4) we recall (2.1) hence we have the following expansion:

$$f_{nT}(x|\mathbf{a}_n) = B_n(\mathbf{a}_n)f_C(x)(\frac{1}{2^n}+\frac{1}{2^n}\sum_{j=0}^{\infty}U_j(x)\sum_{k=1}^{n}b_{n,k}a_k^j).$$

where we denoted $b_{n,k}=\prod_{j=1,j\neq k}^{n}\frac{\gamma(a_k,a_j)}{(a_k-a_j)(1-a_ka_j)}$. Now recall that

$$\int_{-1}^{1}U_j(x)f_C(x)dx=\left\{\begin{array}{lll}0 & if & j\text{ is odd}\\ 1 & if & j\text{ is even}\end{array}\right. .$$

Hence we have relationship $1=B_n(\mathbf{a}_n)(1/2^n+(\sum b_{n.k}/(1-a_k^2))/2^n)$.

To get the second assertion we use expansion (4.5) and identity (4.7). We get then:

$$f_{nT}(x|\mathbf{a}_n) = B_n(\mathbf{a}_n)f_C(x)(\frac{1}{2^n}+\frac{1}{2^n}\sum_{k=1}^{n}b_{n,k}\frac{(1-a_k^2)}{(1+a_k^2-2a_kx)}=$$

$$B_n(\mathbf{a}_n)f_C(x)(\frac{1}{2^n}+\frac{1}{2^{n-1}}\sum_{k=1}^{n}b_{n,k}(\frac{1-a_kx}{(1+a_k^2-2a_kx)}-\frac{1}{2})=$$

$$B_n(\mathbf{a}_n)f_C(x)(\frac{1}{2^n}-\frac{1}{2^n}\sum_{k=1}^{n}b_{n,k}+\frac{1}{2^{n-1}}\sum_{k=1}^{n}b_{n,k}\frac{1-a_kx}{(1+a_k^2-2a_kx)})=$$

$$B_n(\mathbf{a}_n)f_C(x)(\frac{1}{2^{n-1}}-\frac{1}{2^n}-\frac{1}{2^n}\sum_{k=1}^{n}b_{n,k}+\frac{1}{2^{n-1}}\sum_{k=1}^{n}b_{n,k}\frac{1-a_kx}{(1+a_k^2-2a_kx)}).$$

Further recalling expansion (1.3) we get

$$f_{nT}(x|\mathbf{a}_n) = B_n(\mathbf{a}_n)f_C(x)(\frac{1}{2^{n-1}}-\frac{1}{B_n(\mathbf{a}_n)})+\frac{B_n(\mathbf{a}_n)}{2^{n-1}}f_C(x)\sum_{k=1}^{n}b_{n,k}\sum_{j=0}^{\infty}a_k^jT_j(x)$$

$$=(B_n(\mathbf{a}_n)/2^{n-1}-1)f_C(x)+f_C(x)\frac{B_n(\mathbf{a}_n)}{2^{n-1}}\sum_{k=1}^{n}b_{n,k}+\frac{B_n(\mathbf{a}_n)}{2^{n-1}}f_C(x)\sum_{j=1}^{\infty}T_j(x)\sum_{k=1}^{n}b_{n,k}a_k^j$$

$$=(B_n(\mathbf{a}_n)/2^{n-1}-1)f_C(x)+f_C(x)B_n(\mathbf{a}_n)(\frac{2}{B_n(\mathbf{a}_n)}-\frac{1}{2^{n-1}})$$

$$+\frac{B_n(\mathbf{a}_n)}{2^{n-1}}f_C(x)\sum_{j=1}^{\infty}T_j(x)\sum_{k=1}^{n}b_{n,k}a_k^j=f_C(x)+\frac{B_n(\mathbf{a}_n)}{2^{n-1}}f_C(x)\sum_{j=1}^{\infty}T_j(x)\sum_{k=1}^{n}b_{n,k}a_k^j$$

Finally we recall definition of $B_n(\mathbf{a}_n)$. □

REFERENCES

[1] Bressoud, D. M. A simple proof of Mehler's formula for q-Hermite polynomials. *Indiana Univ. Math. J.* **29** (1980), no. **4**, 577–580. MR0578207 (81f:33009)

[2] Ismail, Mourad E. H. Classical and quantum orthogonal polynomials in one variable. With two chapters by Walter Van Assche. With a foreword by Richard A. Askey. Encyclopedia of Mathematics and its Applications, 98. *Cambridge University Press,* Cambridge, 2005. xviii+706 pp. ISBN: 978-0-521-78201-2; 0-521-78201-5 MR2191786 (2007f:33001)

[3] Paweł J Szabłowski, Askey-Wilson integral and its generalizations, *Advances in Difference Equations* 2014, **2014:316**

[4] Szabłowski, Paweł J. On affinity relating two positive measures and the connection coefficients between polynomials orthogonalized by these measures. *Appl. Math. Comput.* **219** (2013), no. 12, 6768–6776. MR3027843
[5] Szabłowski, Paweł. J, On generalized Kesten–McKay distributions, submitted, http://arxiv.org/abs/1507.03191
[6] Szabłowski, Paweł J. Around Poisson–Mehler summation formula, submitted, http://arxiv.org/abs/1108.3024

Department of Mathematics and Information Sciences,, Warsaw University of Technology, ul Koszykowa 75, 00-662 Warsaw, Poland
E-mail address: `pawel.szablowski@gmail.com`